

VIDREL® PreFilled Syringe

(choriogonadotropin alfa injection)

Ovidrel® PreFilled Syringe (choriogonadotropin alfa injection) is the only recombinant, liquid, ready-to-inject human chorionic gonadotropin (r-hCG) approved in the world. It is manufactured using the Recombinant technology, which provides high purity and consistency from batch to batch.

Administration of Ovidrel® PreFilled Syringe

Get ready

1. Make sure you have all the necessary materials assembled in a clean area: Ovidrel® PreFilled Syringe, alcohol swabs, gauze, sharps-disposable container.



Please allow the pre-filled syringe to adjust to room temperature before you administer your injection.

2. Wash your hands thoroughly.



Prepare your Ovidrel® PreFilled Syringe

3. With the needle pointing upwards, carefully remove the needle cap from the syringe. Do not touch the needle or allow it to touch any surface. Keep materials sterile.



4. To remove any air bubbles, point the needle up and gently tap on the syringe until all the bubbles rise to the top.



5. Push the plunger carefully until a small drop of liquid begins to appear from the tip of the needle.



Administration and storage instructions

Prepare the injection area

6. Choose an injection site in the lower abdominal area, preferably around the belly button but at least 1 inch away.



7. Carefully clean the injection site on the stomach with an alcohol swab and allow it to air-dry.



Administer your injection

8. Holding the syringe with one hand the way you would hold a pencil, pinch the skin on the chosen injection site with the other hand and hold firmly.



9. Insert the entire length of the needle into the skin at an upward angle of about 45 to 90 degrees, as indicated by your doctor or nurse.



10. Release the skin and push the plunger in a slow, steady motion until all the medication is injected. Take as much time as you need to inject all the contents.



11. After injecting all the contents, gently withdraw the needle.

12. Apply pressure to the injection site with a gauze pad. If bleeding does not stop within a few minutes, place a piece of clean gauze over the injection site and cover it with an adhesive bandage.



Discard used materials

13. Discard the syringe in your sharps-disposal container. Remember that injection materials must be kept sterile and cannot be reused.



Always take your injection exactly as your doctor has instructed.

Storage of Ovidrel[®] PreFilled Syringe

Important: Ovidrel[®] PreFilled Syringe should be stored refrigerated (36°- 46°F / 2-8°C) to allow the product to be used until the expiry date shown on the syringe or carton. Alternatively, the Ovidrel[®] PreFilled Syringe may be stored by the patient for no more than 30 days at room temperature (up to 77°F / 25°C) and in this case must be used within those 30 days. Store away from the light. Do not freeze.

